

# Diabetes in British Columbia

**2023 Background**

**Summary:** This backgrounder provides key statistics about diabetes in British Columbia, the impact of diabetes on the population of British Columbia, and Diabetes Canada's recommendations to the Government of British Columbia to address diabetes prevention and management.

**Publication Date:** July 2023

**Report Length:** 6 Pages

**Cite As:** Diabetes in British Columbia: Backgrounder. Ottawa: Diabetes Canada; 2023.

**About Diabetes Canada:** Diabetes Canada is a national health charity representing more than 11.9 million people in Canada living with diabetes or prediabetes. Diabetes Canada leads the fight against diabetes by helping those affected by diabetes live healthy lives, preventing the onset and consequences of diabetes, and discovering a cure. It has a heritage of excellence and leadership, and its co-founder, Dr. Charles Best, along with Dr. Frederick Banting, is credited with the co-discovery of insulin. Diabetes Canada is supported in its efforts by a community-based network of volunteers, employees, health care professionals, researchers, and partners. By providing education and services, advocating on behalf of people living with diabetes, supporting research, and translating research into practical applications, Diabetes Canada is delivering on its mission. Diabetes Canada will continue to change the world for those affected by diabetes through healthier communities, exceptional care, and high-impact research.

For more information, please visit: [www.diabetes.ca](http://www.diabetes.ca)

**Contact:** [advocacy@diabetes.ca](mailto:advocacy@diabetes.ca) with inquiries about this Diabetes Canada report.


## Estimated Prevalence and Cost of Diabetes – British Columbia

Prevalence (1)	2023	2033
Diabetes (type 1 + type 2 diagnosed + type 2 undiagnosed)	852,000 / 16%	1,097,000 / 20%
Diabetes (type 1 and type 2 diagnosed)	596,000 / 11%	768,000 / 14%
Diabetes (type 1)	5-10% of diabetes prevalence	
Diabetes (type 1 + type 2 diagnosed + type 2 undiagnosed) and prediabetes (includes undiagnosed)	1,671,000 / 32%	2,000,000 / 36%
Increase in diabetes (type 1 and type 2 diagnosed), 2023-2033	29%	
Direct cost to the health care system	\$583 million	\$747 million
Out-of-pocket cost per year (2)		
Type 1 diabetes costs, % of family income	\$945-\$6,487 / 3%-4%	
Type 2 diabetes costs, % of family income	\$388-\$6,098 / 1%-4%	

### Impact of Diabetes

- Among British Columbians (1):
  - **32%** live with diabetes or prediabetes and
  - **11%** live with diagnosed diabetes, a figure that climbs to **16%** when cases of undiagnosed type 2 diabetes are included.
- Diabetes complications are associated with premature death (3). Diabetes can reduce lifespan by **five to 15 years** (3). It is estimated that the all-cause mortality rate among people living with diabetes is **twice** as high as the all-cause mortality rate for those without diabetes (4).
- People with diabetes are over **three times** more likely to be hospitalized with cardiovascular disease, **12 times** more likely to be hospitalized with end-stage renal disease, and almost **20 times** more likely to be hospitalized for a non-traumatic lower limb amputation compared to the general population (3).

- Diabetes contributes to (5):



- **33-50%** of people living with diabetes experience diabetes distress (an overwhelming feeling about their condition that can lead to unhealthy habits like not checking their blood sugar or skipping medical appointments, etc.) (6).

- Individuals with depression have a **40% – 60%** increased risk of developing type 2 diabetes (6).
  - Diabetic retinopathy, a retinal vascular disorder that occurs as a complication of diabetes, is a leading cause of new cases of blindness in Canada, and often affects working-aged adults (7).
  - Vision loss is associated with significant morbidity, including increased falls, hip fractures, and an increased risk of death (8).
  - Foot ulceration affects an estimated **15-25%** of people with diabetes in their lifetime (9).
  - Compared to the general population, adults living with diabetes in Canada are over **20 times** more likely to undergo non-traumatic lower limb amputations - 85% of which are preceded by foot ulcers (10).
  - Hypoglycemia (low blood sugar) and hyperglycemia (high blood sugar) may affect mood and behaviour and can lead to emergency situations if left untreated (11).
  - For people living with diabetes, adherence to treatment is affected by costs which are not covered by their public drugs and devices coverage (2).
    - Those with type 1 diabetes can pay up to 4% of their gross annual income on medications and devices that range from \$945 to \$6,487.
    - Those living with type 2 diabetes can pay up to 4% of their gross annual income on medications and devices that range from \$388 to \$6,098.
  - Type 2 diabetes is caused by a combination of individual, social, environmental, and genetic factors (11).
  - Certain populations are at higher risk of developing type 2 diabetes, such as those of African, Arab, Asian, Hispanic, Indigenous, or South Asian descent, those who are older, have a lower level of income or education, are physically inactive, or are living with overweight or obesity (11).
  - The age-standardized prevalence rates for diabetes are **16%** among people of South Asian descent, **13.3%** among Black adults, **12.5%** among people of Arab/West Asian descent, **8.8%** among people of East/Southeast Asian descent, and **5.7%** among people of Latin American descent (12).
  - The prevalence of diabetes among First Nations adults living off reserve, Metis adults, and Inuit adults is **1.72 times**, **1.22 times**, and **1.18 times** higher respectively than the prevalence among non-Indigenous adult (12). In addition to the risk factors that impact all people in Canada, the ongoing burden of colonization continues to influence Indigenous peoples' health.
  - The prevalence of diabetes among adults in the lowest income groups is **2.1 times** that of adults in the highest income group (12).
  - Adults who have not completed high school have a diabetes prevalence **1.9 times** that of adults with a university education (12).
  - Social determinants of health can influence the rate of individual-level modifiable risk factors and thus the risk of diabetes. The main determinants of health include income, employment, education, childhood experiences,
- 
- ### Risk Factors for Diabetes
- The risk factors for type 1 diabetes are not well understood, but interaction between genetic and environmental factors are likely involved (11).

physical environments, social supports, access to health services, and racism (13).

---

### Policy, Programs, and Services Related to Diabetes

---

- On August 1, 2023 the government announced coverage of the FreeStyle Libre 2 isCGM according to the rules of B.C. PharmaCare for aged 4+ requiring basal and bolus insulins.
- As of January 1, 2023, all five B.C Health Authorities can provide total contact cast (TCC) services for treatment of diabetic foot ulcers. In addition, for clients with BC Pharmacare coverage, custom orthotics and adaptive shoes will be covered for those with diabetic foot ulcers recently healed with total contact casting.
- In November 2021, the government eliminated its MAC pricing policy for rapid-acting insulins and now covers their full cost.
- In June 2021, the government announced coverage of the Dexcom G6 rtCGM according to the rules of B.C. PharmaCare for aged 2+ requiring basal and bolus insulins.
- Effective April 2021, the government introduced a tax of seven percent on soda beverages to help address obesity and type 2 diabetes.
- On January 1, 2020, Medical Service Plan (MSP) premiums were eliminated for all British Columbians.
- In May 2019, the government launched its Biosimilars Initiative which introduced changes for children and adults using insulin. To maintain public coverage, Individuals using an originator insulin will be required to switch to its biosimilar.
- In November 2019, the government announced its commitment to using the

Diabetes 360° framework to develop a provincial diabetes strategy.

- On January 1, 2019, the government eliminated PharmaCare deductibles for households earning up to \$30,000 in annual net income and reduced deductibles for households earning up to \$45,000.
- In July 2018, the government eliminated age restrictions from its insulin pump program. At the same time, the program was tiered, providing one pump model at no charge and others available through PharmaCare if special authority approval is granted.
- In 2018 Health Quality BC partnered with the Institute for Healthy System Transformation and Sustainability (IHSTS) to lead the formation of the Type 2 Diabetes Network focused on sharing, developing and applying innovative practices to improve outcomes for British Columbians living with type 2 diabetes.
- In 2014, the government introduced provincial standards to support students with diabetes in schools, which includes staff training to administer glucagon in an emergency, as well as monitor blood sugar and administer insulin for students who are unable to manage their diabetes independently.

---

### Challenges

---

British Columbia faces unique challenges in preventing type 2 diabetes and meeting the needs of those living with diabetes:

- Non-modifiable risk factors of type 2 diabetes include age, gender, and ethnicity (11).
  - The median age in British Columbia is **43 years** (14). **18.3%** of British Columbians are over 65 years old (14). The risk of developing type 2 diabetes

- increases with age (11). Older adults living with diabetes are more likely to be frail and progressive frailty has been associated with reduced function and increased mortality (15).
- Adult men are more at risk of type 2 diabetes compared to adult women (11).
  - Approximately **31.6%** of British Columbians self-identify as being of African, Arab, Asian, Hispanic, or South Asian descent (14). These groups are at increased risk of developing type 2 diabetes (11).
  - There are **270,585** Indigenous people in British Columbia, who face significantly higher rates of diabetes and adverse health consequences than the overall population (16).
  - British Columbia has high rates of individual-level modifiable risk factors (17):
    - **34.5%** of adults and **74.1** of youth (aged 12 to 17 years) are physically inactive;
    - **35.2%** of adults are living with overweight and **22.2%** of adults are living with obesity;
    - **70.4%** of adults are not eating enough fruits and vegetables; and
    - **14.1%** of adults are current tobacco smokers.
  - Factors related to the social determinants of health and that can influence the rate of individual-level modifiable risk factors among British Columbians include income, education, food security, the built environment, social support, and access to health care (3).

---

## Diabetes Canada's Recommendations to the Government of British Columbia

---

- 1. Fund a provincial diabetes framework**
  - Allocate funding to develop and implement a comprehensive diabetes framework with measurable goals for improving diabetes prevention, treatment and health outcomes for the province.
- 2. Expand access: Put patients at the centre of policy decisions**
  - Eliminate barriers to access evidence-based, personalized diabetes treatments, including diabetes medications, devices, and supplies.
  - Provide personal choice and equitable access to insulin pumps and continuous glucose monitoring systems (isCGM & rtCGM) according to Diabetes Canada's CGM [reimbursement recommendations](#).
- 3. Prevent amputations**
  - Expand health policies that support the prevention and management of diabetes foot complications and reduce the risk of lower limb amputations.

---

## References

---

1. Canadian Diabetes Cost Model. Ottawa: Diabetes Canada; 2016.
2. Diabetes and Diabetes-Related Out-of-Pocket Costs: 2022 Update. Diabetes Canada; 2022.
3. Diabetes in Canada: Facts and figures from a public health perspective [Internet]. Ottawa: Public Health Agency of Canada; 2011 p. 126. Available from:

- <https://www.canada.ca/content/dam/phac-aspc/migration/phac-aspc/cd-mc/publications/diabetes-diabete/facts-figures-faits-chiffres-2011/pdf/facts-figures-faits-chiffres-eng.pdf>
4. Public Health Agency of Canada. Twenty years of diabetes surveillance using the Canadian chronic disease surveillance system. [Internet]. 2019. Available from: [http://publications.gc.ca/collections/collectio n\\_2019/aspc-phac/HP35-122-2019-eng.pdf](http://publications.gc.ca/collections/collectio n_2019/aspc-phac/HP35-122-2019-eng.pdf)
  5. Hux J, Booth J, Slaughter P, Laupacis A. Diabetes in Ontario: An ICES Practice Atlas [Internet]. Institute for Clinical Evaluative Sciences; 2003. Available from: <https://www.ices.on.ca/Publications/Atlases-and-Reports/2003/Diabetes-in-Ontario>
  6. Diabetes Canada Clinical Practice Guidelines Expert Committee, Robinson DJ, Coons M, Haensel H, Vallis M, Yale JF. Diabetes and Mental Health. *Can J Diabetes*. 2018 Apr;42 Suppl 1:S130–41.
  7. Flaxel, Christina J.; Adelman, Ron A.; Bailey, Steven T.; Fawzi, Amani; Lim, Jennifer I.; Vemulakonda, G. Atma; Ying, Gui-shuang. *Ophthalmology (Rochester, Minn.)*, 2020, Vol.127 (1), p.P66-P145
  8. 10. Vu, H. T., Keeffe, J. E., McCarty, C. A., & Taylor, H. R. (2005). Impact of unilateral and bilateral vision loss on quality of life. *The British journal of ophthalmology*, 89(3), 360–363. <https://doi.org/10.1136/bjo.2004.047498>
  9. Yazdanpanah, L., Shahbazian, H., Nazari, I., Arti, H. R., Ahmadi, F., Mohammadianinejad, S. E., Cheraghian, B., & Hesam, S. (2018). Incidence and Risk Factors of Diabetic Foot Ulcer: A Population-Based Diabetic Foot Cohort (ADFC Study)-Two-Year Follow-Up Study. *International journal of endocrinology*, 2018, 7631659. <https://doi.org/10.1155/2018/7631659>
  10. Armstrong DG, Boulton AJM, Bus SA. Diabetic Foot Ulcers and Their Recurrence. *N Engl J Med*. 2017 Jun 15;376(24):2367–75
  11. Diabetes Canada Clinical Practice Guidelines Expert Committee. Diabetes Canada 2018 Clinical Practice Guidelines for the Prevention and Management of Diabetes in Canada. *Can J Diabetes*. 2018;42(Suppl 1):S1:S325.
  12. Public Health Agency of Canada. Pan-Canadian Health Inequalities Data Tool, 2022 Edition [Internet]. Available from: <https://health-infobase.canada.ca/health-inequalities/data-tool/>
  13. Prinjha S, Wicklow B, Nakhla M, Banerjee AT. Toward the Goal of Understanding and Tackling the Social Determinants of Diabetes. *Can J Diabetes*. 2022 Aug 1;46(6):549–50.
  14. Government of Canada SC. Census Profile, 2016 Census -British Columbia [Province] and Canada [Country] [Internet].
  15. Meneilly GS, Knip A, Miller DB, Sherifali D, Tessier D, Zahedi A. Diabetes in Older People. *Can J Diabetes*. 2018 Apr;42: S283–95.
  16. Aboriginal peoples in Canada: Key results from the 2016 Census [Internet]. Ottawa: Statistics Canada; 2017 Oct p. 11. Available from: [https://www150.statcan.gc.ca/n1/en/daily-quotidien/171025/dq171025a-eng.pdf?st=krvs\\_yCt](https://www150.statcan.gc.ca/n1/en/daily-quotidien/171025/dq171025a-eng.pdf?st=krvs_yCt)
  17. *Canadian Risk Factor Atlas (CRFA), 2020 edition*. Public Health Agency of Canada. Available at Public Health

## Diabetes in British Columbia

Infobase: <https://health-infobase.canada.ca/crfa/>